

Bryant University

Bryant Digital Repository

Science and Technology Faculty Conference
Proceedings

Science and Technology Faculty Publications
and Research

2018

Interpreting long chain n-alkyl lipids in lake sediments: A critical evaluation of their sources and isotope signals

Hong Yang

Bryant University, hyang@bryant.edu

Qin Leng

Bryant University, qleng@bryant.edu

Katelynn Howard

Kai-Lou Yue

Follow this and additional works at: <https://digitalcommons.bryant.edu/sci-conf-proceedings>



Part of the [Earth Sciences Commons](#), [Environmental Sciences Commons](#), and the [Life Sciences Commons](#)

Recommended Citation

Yang, Hong; Leng, Qin; Howard, Katelynn; and Yue, Kai-Lou, "Interpreting long chain n-alkyl lipids in lake sediments: A critical evaluation of their sources and isotope signals" (2018). *Science and Technology Faculty Conference Proceedings*. Paper 2.

<https://digitalcommons.bryant.edu/sci-conf-proceedings/2>

This Conference Proceeding is brought to you for free and open access by the Science and Technology Faculty Publications and Research at Bryant Digital Repository. It has been accepted for inclusion in Science and Technology Faculty Conference Proceedings by an authorized administrator of Bryant Digital Repository. For more information, please contact dcommons@bryant.edu.

Interpreting long chain *n*-alkyl lipids in lake sediments: A critical evaluation of their sources and isotope signals

HONG YANG*, QIN LENG, KATELYNN HOWARD, KAILOU YUE

Laboratory for Terrestrial Environments, College of Arts and Sciences, Bryant University, Smithfield, Rhode Island 02917, USA; *correspondence: hyang@bryant.edu

Stable carbon and hydrogen isotopic values of *n*-alkyl lipids extracted from lake sediments have been widely used for the reconstruction of ancient vegetation, climate, altitude, hydrology, etc. Compound-specific isotope analysis has taken the advantage of source-specific environmental signals from both autochthonous and allochthonous lipids, assuming that long chain *n*-alkyl lipids are primary derived from terrestrial plants. In recent years, substantial progresses have been made toward lipids produced by lacustrine macrophytes (aquatic plants and macro-algae) regarding their abundance, metabolism, chain length distribution, and isotope values.

Accumulating evidence indicates that long chain (C_{26} – C_{31}) *n*-alkyl lipids in lacustrine sediments are derived from diverse sources, including terrestrial plants, aquatic macrophytes, and microbes; some emerged aquatic plants and macro-algae produce similar or even higher concentrations of long chain *n*-alkyl lipids as they are from terrestrial plants. Abundant long chain *n*-alkyl lipids from submerged aquatic plants with C_4 -like metabolism suggested an underlying cause for the large discrepancies of $\delta^{13}C$ values and unformed δD value variations among homologous lipids with different chain lengths, complicating the estimate of ancient C_3/C_4 vegetation and the reconstruction of paleohydrology. The significant correlations between δD values of long-chain *n*-alkyl lipids from aquatic macrophytes and lake water (*v/s* δD values of precipitation) found in lakes at the Tibetan Plateau further highlight both challenges and opportunities for paleohydrological reconstructions.

The assessment of relative contributions of long-chain *n*-alkyl lipids from various sources in a particular lacustrine site becomes a critical prerequisite for the application of their isotope values as proxies. To discriminate the sources of these lipids, we found that some traditional indicators, such as Paq, ACL, and CPI, are of limited values, while inter-molecular isotope offsets among *n*-alkyl lipids with different chain lengths show promising results. Knowledge gaps exist regarding factors that influence lipid biomass from aquatic macrophytes, their molecular isotope behaviors, and implications as lacustrine environmental proxies.